

## LETTER

# LightCycler SeptiFast technology in patients with solid malignancies: clinical utility for rapid etiologic diagnosis of sepsis

Lenka Dubská<sup>1\*</sup>, Martina Vyskočilová<sup>1</sup>, Dagmar Minaříková<sup>2</sup>, Petr Jelínek<sup>2</sup>, Renata Tejkalová<sup>3</sup> and Dalibor Valík<sup>1</sup>

**[AU Query: Very carefully check and confirm meaning of text remains OK throughout]**

Recent development of molecular tools for pathogen detection introduced the possibility of early targeted antimicrobial treatment that, in turn, may improve the outcome of patients with sepsis [1].

We retrospectively evaluated 54 results of the SeptiFast test from patients with solid malignancy admitted to the ICU between June 2009 and August 2011. Specimens from suspected bloodstream infection were analyzed using LightCycler SeptiFast (Roche Molecular Diagnostics [AU Query: Provide city, state and country]) according to the manufacturer's instruction and evaluated in comparison with blood culture results obtained from blood sampled no longer than 24 hours before or after sampling for SeptiFast. Blood culturing and identification were performed according to the routine diagnostic procedures. The total number of blood cultures analyzed was 85, and finding a microorganism by either of the two tests was evaluated as positive [AU Query: Confirm sentence is OK]. Consistently, negative results from SeptiFast and blood culture were obtained in 21 (39%) cases.

To assess the true positivity of both microbiological methods, discrepant cases were evaluated in the context of clinical and laboratory findings by two independent physicians experienced in critical care (Figure 1). The clinically relevant presence of a pathogen detected by blood culture but not by SeptiFast was recorded for *Klebsiella pneumoniae/oxytoca*, and in both cases the presence of another member of Enterobacteriaceae was reported by SeptiFast, *Escherichia coli* and *Enterobacter cloacae/aerogenes* – thus misidentification of strains with atypical phenotype cannot be excluded [3] [AU Query:

Confirm sentence is OK]. Our results also show that SeptiFast is more efficient in detection of clinically relevant infection by *E. coli* and *Pseudomonas aeruginosa*.

The turnaround time for blood culture analysis is 24 to 48 hours. The workflow in our laboratory allows SeptiFast analysis every working day during a day shift. In such conditions, the turnaround time of SeptiFast results for samples delivered to the laboratory on a workday between 7:00 and 14:00 hours was as follows: 100% samples, 10 hours; 97%, 8 hours; 59%, 6 hours [AU Query: Confirm sentence is OK]. The average turnaround time for tests performed with manual DNA isolation was 6 hours 11 minutes; by implementation of automated isolation of DNA using the MagNA Pure Compact System [4], the mean turnaround time was shortened to 5 hours 17 minutes [AU Query: Confirm sentence is OK].

In conclusion, detection of pathogen by the LightCycler SeptiFast system is generally not inferior to the results from blood culture. The added value of multiplex DNA amplification-based pathogen detection is the shorter turnaround time and probably the increased true sensitivity of detection of certain pathogens [AU Query: Confirm sentence is OK]. Moreover, the subset of pathogen detection is required in the background of antibiotic administration; in such conditions, cultivation-independent methods are of clear benefit [5] [AU Query: Confirm sentence is OK].

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

LD, MV, RT and DV participated in the design of the study. LD and MV carried out the molecular genetics part of the study. DM and PJ evaluated the microbiological findings in the context of clinical status. All authors participated in writing the report.


## Acknowledgements

The authors received a contribution for the reagents, equipment from Roche Diagnostics for this project, and were supported by the European Regional Development Fund and the State budget of the Czech Republic for RECAMO (CZ.1.05/2.1.00/03.0101) [AU Query: Confirm sentence is OK].

\*Correspondence: [dubska@mou.cz](mailto:dubska@mou.cz)

<sup>1</sup>Department of Laboratory Medicine, Masaryk Memorial Cancer Institute, Zluty kopec 7, 656 53 Brno, Czech Republic

Full list of author information is available at the end of the article



#### Author details

<sup>1</sup>Department of Laboratory Medicine, Masaryk Memorial Cancer Institute, Zluty kopec 7, 656 53 Brno, Czech Republic. <sup>2</sup>Department of Anesthesiology and Intensive Care, Masaryk Memorial Cancer Institute, Zluty kopec 7, 656 53 Brno, Czech Republic. <sup>3</sup>Department of Microbiology, St. Anne's Faculty Hospital and Faculty of Medicine, Masaryk University, Pekarska 53, 664/53, 656 91 Brno, Czech Republic.

Published: X Month 2012

#### References

1. Garnacho-Montero J, Ortiz-Leyba C, Herrera-Melero I, Aldabó-Pallás T, Cayuela-Dominguez A, Marquez-Vacaro JA, Carbajal-Guerrero J, Garcia-Garmendia JL: **Mortality and morbidity attributable to inadequate empirical antimicrobial therapy in patients admitted to the ICU with sepsis: a matched cohort study.** *J Antimicrob Chemother* 2008, **61**:436-441.
2. Roche Molecular Diagnostics [http://molecular.roche.com/assays/Pages/LightCyclerSeptiFastTestMGRADE.aspx]
3. Claes G, De Baere T, Wauters G, Vandecandelaere P, Verschraegen G, Muylaert A, Vanechoutte M: **Extended-spectrum beta-lactamase (ESBL) producing**

*Enterobacter aerogenes* phenotypically misidentified as *Klebsiella pneumoniae* or *K. terrigena*. *BMC Microbiol* 2004, **4**:49.

4. Regueiro BJ, Varela-Ledo E, Martinez-Lamas L, Rodriguez-Calviño J, Aguilera A, Santos A, Gomez-Tato A, Alvarez-Escudero J: **Automated extraction improves multiplex molecular detection of infection in septic patients.** *PLoS One* 2010 **5**:e13387.
5. Yanagihara K, Kitagawa Y, Tomonaga M, Tsukasaki K, Kohno S, Seki M, Sugimoto H, Shimazu T, Tasaki O, Matsushima A, Ikeda Y, Okamoto S, Aikawa N, Hori S, Obara H, Ishizaka A, Hasegawa N, Takeda J, Kamiyama S, Sugahara K, Asari S, Murata M, Kobayashi Y, Ginba H, Sumiyama Y, Kitajima M: **Evaluation of pathogen detection from clinical samples by real-time polymerase chain reaction using a sepsis pathogen DNA detection kit.** *Crit Care* 2010, **4**:R159.

doi:10.1186/cc10595

**Cite this article as:** Dubská L, et al.: LightCycler SeptiFast technology in patients with solid malignancies: clinical utility for rapid etiologic diagnosis of sepsis. *Critical Care* 2012, **16**:4??.